



Travis G. Maak, M.D.
 590 Wakara Way
 Salt Lake City, UT 84108
 Tel: (801) 587-7040
 Fax: (801)587-7112
 Lic. # 8234797-1205

Physical Therapy Prescription Microfracture

Patient Name: _____

Today's Date: _____

Surgery Date: _____

Dx: s/p (LEFT / RIGHT) Knee Microfracture - (MFC / LFC / TROCHLEA / PATELLA / MTP / LTP)

Modalities:

Week 0 - 4

- ___ Ice / Massage / Anti-Inflammatory Modalities
- ___ Ambulate NWB / TTWB / PWB / FWB in Bledsoe Brace locked @ 0°
- ___ Limit ROM to 30° for 4 weeks
- ___ Range of Motion Active / Active-Assisted / Passive
- ___ Home CPM 1-2 hrs / session, TID for 6 weeks
- ___ LE musculature mobilization and stretching
- ___ Gradual Progression of Resistance Exercises in Flexed Knee, OKC only
- ___ Quadriceps Strengthening
 - ___ 0-30° Arc
- ___ Hamstring, Gluteal, and lateral hip strengthening
- ___ Begin Straight Leg Raises (Knee at 0°)
- ___ Quad Isometrics
- ___ UBE for cardiovascular training
- ___ Heel chord stretching
- ___ Electrical Stimulation for Quadriceps
- ___ Hydrotherapy once incisions are fully closed

Week 4-6

- ___ Begin to progress to full ROM. No restrictions on motion, limit CKC to 90 degrees of flexion
- ___ May begin to WBAT in brace, limit motion to 30 degrees of flexion when ambulating
- ___ May begin proprioception exercises on stable surface
- ___ Core and hip strength and endurance

Week 6-12

- ___ May discontinue use of brace. No motion restrictions in OKC or CKC
- ___ Begin full arc CKC strength focused on quadriceps, hamstring and gluteal muscle groups
- ___ Progress proprioception to unstable surface with perturbations
- ___ Assess posture and functional movement patterns. Corrective exercise as needed
- ___ Progress core and hip strength and endurance
- ___ Okay to begin frontal plane CKC strength, avoid pivoting
- ___ Continue to work on joint mobility, single and multiple

Week 12+

- ___ Continue CKC strength
- ___ Begin light agility drills in frontal and sagittal plane only, no pivoting until 4.5 – 5 months post op
- ___ Progress to multi planar core strengthening
- ___ May begin in line jogging if eccentric step down is symmetric
- ___ Okay to begin Olympic lifting and triple extension exercises at 4.5 months post op

Frequency & Duration: (circle one) 1-2 2-3 x/week for _____ weeks Home Program

**Please send progress notes.

Physician's Signature: _____ **M.D.**